



## IUCN Red List Species Richness

These EnviroAtlas national maps display the number of terrestrial vertebrate species listed as threatened by the International Union for Conservation of Nature (IUCN, version 2016-1) with potential habitat within each 12-digit hydrologic unit ([HUC](#)) in the conterminous United States. These data are based on habitat models, not wildlife counts.

### Why are IUCN-listed species important?

The metric, IUCN Red List Species Richness, estimates the species abundance of terrestrial vertebrate species listed as threatened based on potential habitat.<sup>1</sup> The IUCN assesses conservation status of species at a global scale to identify taxa threatened with extinction. The animals assessed for the IUCN Red List are important for genetic and ecosystem diversity. Information on their conservation status and distribution helps to inform management about conserving biodiversity from local to global scales. The IUCN Red List of Threatened Species provides conservation status and distribution information on animals that have been globally evaluated using the IUCN Red List Categories and Criteria.<sup>1</sup> This system determines the relative risk of extinction, listing species as critically endangered, endangered, and vulnerable. The IUCN Red List also includes animals that would meet the definition of threatened if not for conservation efforts (i.e., they are near-threatened).

Species richness is one measure of [biodiversity](#) that can represent the relative conservation value of a particular area. Many scientists believe that biodiversity, because it represents all forms of life on earth, provides or supports the core benefits that humans derive from their environment to sustain human society, economy, health, and well-being. Managing for biodiversity is one way to balance competing demands for ecosystem services.<sup>2</sup>

Each species plays an important role within its [ecosystem](#), and ecosystems are highly interconnected. Each species depends on others for some aspect of its survival to provide food, habitat, decomposition, pollination, or control of pest species. The removal of even one species from an ecosystem can create a [trophic cascade](#) that can affect the entire [food chain](#). For example, grazers and browsers directly modify the species composition, diversity, and condition of grassland and forest habitats. Top predators, by regulating herbivore numbers, indirectly influence habitat condition and diversity by reducing grazing pressure on plant production.<sup>3</sup>



Characteristics of biodiversity are valued in a variety of ways and thus they are important to include in any assessment that seeks to identify and quantify the value of ecosystems to humans. Some biodiversity metrics clearly reflect ecosystem services and their contribution to our quality of life and economy (e.g., abundance and diversity of game species), whereas others reflect indirect and difficult to quantify relationships to services (e.g., relevance of species diversity to ecosystem resilience, cultural value, and aesthetic amenities).

### How can I use this information?

Three EnviroAtlas maps, Mean, Maximum, and Normalized Index of Biodiversity (NIB) illustrate IUCN Red List Species Richness for terrestrial vertebrate species listed as threatened for each 12-digit HUC in the conterminous United States.<sup>4</sup> Used together or independently, these maps can help identify areas of low or high potential threatened species richness to help inform decisions about resource restoration, use, and conservation. Mean richness is a commonly used and understood value for comparison. NIB provides an index to compare a metric with other metrics across multiple project scales simultaneously. Maximum richness identifies areas that are species rich but may not occupy large areas (e.g. linear riparian areas).

These maps can be used in conjunction with other maps in EnviroAtlas such as ecoregions, the U.S. Geological Survey (USGS) protected areas database ([PAD-US](#)), or the USGS Gap Analysis Project ([GAP](#)) ecological systems to identify

areas with high ecological or recreational value for conservation, recreation, or restoration planning.

After learning the IUCN Red List species richness values for a particular 12-digit HUC, users can investigate an area more intensively by using individual species models available from the GAP Project.

### How were the data for this map created?

The USGS GAP project maps the distribution of natural vegetation communities and potential habitat for individual terrestrial vertebrate species. These models use environmental variables (e.g., land cover, elevation, and distance to water) to predict habitat for each species. GAP modeled habitat for 1,590 terrestrial vertebrate species that reside, breed, or use the habitat within the conterminous U.S. for a significant portion of their life history.

Predicted habitat for the 57 individual IUCN-listed species (Version 2016-1) was combined to calculate Red List species richness by pixel. Of the 57 species, there were 11 amphibian, 7 bird, 27 mammal and 12 reptile species. The mean and maximum numbers of terrestrial vertebrate species in each 30-meter pixel were calculated for each 12-digit HUC. The mean species richness value by HUC was divided by the maximum mean value within all HUCs to calculate the NIB.

### What are the limitations of these data?

EnviroAtlas uses the best data available, but there are still limitations associated with these data. These data, based on models and large national geospatial databases of predicted habitat, are estimations of reality that may overestimate actual species presence. Modeled data are intended to complement rather than replace monitoring data. Habitat models do not predict the actual occurrence of species, but rather their potential occurrence based on their known associations with

certain habitat types. Habitat is only one factor that determines the actual presence of a species. Other factors include habitat quality, predators, prey, competing species, and fine scale habitat features. Other essential species information in addition to species richness includes the types of species and their [functional groups](#), whether they are rare or common, native or non-native, tolerant or intolerant of disturbance.

### How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded. Individual 30-meter pixel data may be downloaded from the [New Mexico State University Center for Applied Spatial Ecology](#).

### Where can I get more information?

A selection of resources related to terrestrial vertebrates and biodiversity is listed below. Information on the models and data used in the USGS Core Science Analytics, Synthesis & Library's [GAP](#) project is available on their website. For additional information on how the data were created, access the [metadata](#) for the data layer from the layer list drop down menu. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

### Acknowledgments

The data for IUCN Red List Species Richness were created through a collaborative effort between the EPA and USGS GAP. GAP habitat models were created by New Mexico State University, North Carolina State University, and Boise State University personnel. Kenneth Boykin and graduate students from New Mexico State University generated the biodiversity metrics. The fact sheet was written by Kenneth Boykin (New Mexico State University), William Kepner, Anne Neale, and Jessica Daniel (EPA), and Megan Culler (EPA Student Services Contractor).

### Selected Publications

1. IUCN 2020. [The IUCN Red List of Threatened Species](#). Version 2020-1. Accessed April 2020.
2. Boykin, K.G., W.G. Kepner, D.F. Bradford, R.K. Guy, D.A. Kopp, A. Leimer, E. Samson, F. East, A. Neale, and K. Gergely. 2013. [A national approach for mapping and quantifying habitat-based biodiversity metrics across multiple spatial scales](#). *Ecological Indicators* 33:139–147.
3. Miller, B., B. Dugelby, D. Foreman, C. Martinez del Rio, R. Noss, M. Phillips, R. Reading, M. E. Soulé, J. Terborgh, and L. Wilcox. 2001. [The importance of large carnivores to healthy ecosystems](#). *Endangered Species Update* 18(5):202–210.
4. Kepner, W.G., K.G. Boykin, D.F. Bradford, A.C. Neale, A.K. Leimer, and K.J. Gergely. 2011. [Biodiversity metrics fact sheet](#), EPA/600/F-11/006, U.S. Environmental Protection Agency, Washington, D.C.
- Hoffmann, M., T.M. Brooks, G.A.B. da Fonseca, C. Gascon, A.F.A. Hawkins, R.E. James, P. Langhammer, R.A. Mittermeier, J.D. Pilgrim, A.S.L. Rodrigues, and J.M.C. Silva. 2008. [Conservation planning and the IUCN Red List](#). *Endangered Species Research* 6:113–125.